

Korn		Prothioconazol 250 g/l		Tebuconazol 200 g/l		Metrafenon 300 g/l	Proquinazid 200 g/l	Pyraclostrobin 200 g/l	Folpet 500 g/l	Pyraclostrobin 100 g/l Meftrifluconazol 100 g/l	Tebuconazol 125 g/l Prothioconazol 125 g/l	Tebuconazol 160 g/l Prothioconazol 80 g/l	Tebuconazol 80 g/l Prothioconazol 160 g/l	Boscalid 500 g/l	Difencconazol 125 g/l Azoxystrobin 125 g/l	Prothioconazol 125 g/l Fluopyram 125 g/l (SDHI)	Prothioconazol 160 g/l Spiroxamin 300 g/l (SDHI)						
Effekt af Fungicider 2022		Proline EC 250		Ortus max 200 EW		Flekty	Tallus	Comet Pro	Folpan 500 SC	Baylay	Prosaro EC 250	Follicur Expert	Proline Xpert	Entargo	Amistar Gold	PROPULSE	Input EC460						
Hvede	kg-/ha	0,40	0,20	0,50	0,25	0,20	0,15	0,25	0,75	1,50	0,75	0,40	0,20	0,40	0,20	0,40	0,20	0,70	0,35	0,50	0,50	0,50	
Forbrug af "triazol-kvote" i %	BBCH 30-32			200%	100%					150%	75%	100%	50%	128%	64%	64%	32%			x			
Triazol bejdset udsæd	BBCH 33-37			100%	50%					150%	75%	50%	25%	64%	32%	32%	16%						
	BBCH 38-69 (40)			48%	24%					100%	50%	24%	12%	30%	15%	15%	8%						
Forbrug af "triazol-kvote" i %	BBCH 30-32			149%	75%					150%	75%	75%	37%	96%	48%	48%	24%			x			
Ubejdset udsæd	BBCH 33-37			75%	38%					150%	75%	38%	19%	48%	24%	24%	12%						
eller Celest Formula M	BBCH 38-69 (40)			40%	20%					100%	50%	20%	10%	26%	13%	13%	6%						
"Tebumidler" med "gammel" etiketter				40%	20%									26%	13%	13%	6%						
		---- max 2 gange pr vækstsæson ----				*1	*1	max 2		*1		-- max 2 gange --						max 1		max 1		max 2	
Erysiphe graminis	Meldug forebyggende	**(*)	**	**	*	**(*)	***(*)		(*)	**	*	**(*)	*	**(*)	*	**(*)	*					***	***
	Meldug kurativ	**		*		*	**					*	(*)	*	(*)	*	(*)					**	**
Puccinia striiformis	Gulrust	**	*(*)	***	**			**	(*)	***	**	***	**	***	**	***	**	*			***	**	***
Puccinia recondita	Brunrust	**	*(*)	***	**			***	(*)	***1	**	***	**	***	**	***	**	*			***	**	**(*)
Septoria tritici	Gråplet	**(*)	**	*					*	****	***	**	*	*(*)	*	**(*)	*	**	*	*	*(*)	***	**
Septoria nodorum	Brunplet	**(*)	**						*	****	***	**	*	*(*)	*	**(*)	*	**	*	*	*(*)	***	**(*)
Drechslera tritici-repentis	Bladplet DTR	***	**							*	**	*	*	*	**	**	*					***	**
Fusarium Spp.	Fusarium	**	*	(*)						0	0	**	*	*	**	*	*					**	*
Pseudocercospora herpotrichoides	Knækkefodsyge	*				**									*		*	**	*		*	*	*
Triazol Bejdset sâsæd	Byg																						
Tebuconazol	BBCH 30-32			187%	93%																		
	BBCH 33-48			118%	59%																		
	BBCH 48-69			78%	39%																		
Mefintri fluconazol	BBCH 33-37									150%	75%												
	BBCH 38-40									100%	50%												
Erysiphe graminis	Bygmeldug	***	**			****	X	*		**	*	**(*)	*	***	**	***	**			X		***	***
Drechslera teres	Bygbladplet	**	*	*(*)	(*)	X	X	***		***	**	*(*)	*	*(*)	*	**	*	**	*	X	X	***	**
Puccinia hordei	Bygrust	***	**	***	**	X	X	****	*	****	***	***	**	***	**	***	**	*		X	X	**	**
Ramularia collo-cygni	Ramularia	***	**			X	X	****		***	**	*	*	*	***	***	**	*		X	X	***	*
Rhynchosporium secalis	Skoldplet	***	**	*(*)	(*)	X	X	**(*)		**	*	***	*	***	**	***	***			X	X	***	***
Kr/ha pr behandling		172	86	83	41	118	98	81	120	713	356	128	64	106	53	144	72	298	149	150	178	215	
Kr/kg/l 2022 listepriis		430		165		590	655	325	160	475	320	265		360		425		300		355		430	

\*\*\*\* = 90-100 % effekt, \*\*\* = 75-90 % effekt, \*\* = 50-75 % effekt, \* < 50 % effekt

\*1: anbefales af hensyn til risiko for resistent kun 1 gang metrafenon pr

d. 19. april 2022 Jørgen Evald Jensen  
planterådgiver  
mail@agriconsult.dk